

Table 1.23  
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites<sup>a</sup> by Race/Ethnicity  
Both Sexes

All Races			White			Black		
	Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>
	2003-2007	1998-2007		2003-2007	1998-2007		2003-2007	1998-2007
All Sites	461.6	-0.9*	All Sites	469.2	-0.9*	All Sites	489.8	-1.4*
Prostate <sup>f</sup>	69.9	-1.2*	Breast	67.9	-2.2*	Prostate <sup>f</sup>	97.8	-2.8*
Breast	66.5	-1.9*	Prostate <sup>f</sup>	67.6	-1.3*	Lung and Bronchus	73.5	-2.2*
Lung and Bronchus	62.5	-1.5*	Lung and Bronchus	63.8	-1.5*	Breast	67.8	-0.4
Colon and Rectum	47.9	-2.6*	Colon and Rectum	47.4	-2.8*	Colon and Rectum	58.9	-1.8*
Urinary Bladder	21.1	-0.4*	Melanoma of the Skin	23.5	2.0*	Pancreas	15.5	0.3
Melanoma of the Skin	20.1	1.6*	Urinary Bladder	22.8	-0.4*	Kidney and Renal Pelvis	15.4	2.1*
Non-Hodgkin Lymphoma	19.6	0.1	Non-Hodgkin Lymphoma	20.5	0.2	Non-Hodgkin Lymphoma	14.7	0.4
Kidney and Renal Pelvis	14.1	2.4*	Kidney and Renal Pelvis	14.5	2.4*	Urinary Bladder	12.8	0.0
Corpus and Uterus, NOS <sup>f</sup>	12.6	-0.7*	Corpus and Uterus, NOS <sup>f</sup>	13.0	-1.0*	Stomach	11.9	-2.6*
Leukemia	12.3	-0.8*	Leukemia	12.9	-0.7*	Corpus and Uterus, NOS <sup>f</sup>	11.8	1.7
Pancreas	11.7	0.7*	Pancreas	11.6	0.8*	Myeloma	11.7	-0.3
Oral Cavity and Pharynx	10.4	-1.1*	Thyroid	10.7	6.3*	Oral Cavity and Pharynx	10.2	-3.1*
Thyroid	10.2	6.1*	Oral Cavity and Pharynx	10.5	-0.8*	Leukemia	9.8	-1.0
Stomach	7.8	-1.7*	Ovary <sup>fh</sup>	7.3	-2.0*	Liver & IBD <sup>g</sup>	8.6	3.6*
Ovary <sup>fh</sup>	7.0	-1.8*	Brain and ONS <sup>g</sup>	7.1	-0.6	Thyroid	6.2	6.7*

  

Asian/Pacific Islander			American Indian/Alaska Native <sup>d</sup>			Hispanic <sup>e</sup>		
	Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>		Rate <sup>b</sup>	APC <sup>c</sup>
	2003-2007	1998-2007		2003-2007	1998-2007		2003-2007	1998-2007
All Sites	310.8	-0.7*	All Sites	317.7	-0.6	All Sites	342.5	-1.3*
Breast	49.1	-0.4	Lung and Bronchus	45.3	0.8	Prostate <sup>f</sup>	55.2	-2.2*
Colon and Rectum	39.2	-2.1*	Colon and Rectum	42.1	-1.7	Breast	46.0	-1.5*
Prostate <sup>f</sup>	39.0	-1.2	Breast	41.4	0.3	Colon and Rectum	37.3	-1.7*
Lung and Bronchus	38.7	-1.0*	Prostate <sup>f</sup>	34.4	-0.7	Lung and Bronchus	32.1	-2.3*
Liver & IBD <sup>g</sup>	14.4	0.6	Kidney and Renal Pelvis	17.4	3.8*	Non-Hodgkin Lymphoma	16.5	-0.2
Stomach	13.2	-3.1*	Non-Hodgkin Lymphoma	11.4	-0.4	Kidney and Renal Pelvis	13.3	1.8*
Non-Hodgkin Lymphoma	12.9	-1.0	Stomach	10.9	-3.7	Stomach	11.6	-2.0*
Thyroid	10.1	4.4*	Liver & IBD <sup>g</sup>	10.6	0.8	Urinary Bladder	11.0	-0.5
Corpus and Uterus, NOS <sup>f</sup>	9.4	1.0	Pancreas	10.3	1.1	Pancreas	10.5	0.5
Urinary Bladder	9.1	-0.2	Corpus and Uterus, NOS <sup>f</sup>	9.0	0.4	Liver & IBD <sup>g</sup>	10.5	1.7*
Pancreas	9.1	0.6	Urinary Bladder	7.7	-	Leukemia	9.5	-0.6
Oral Cavity and Pharynx	7.6	-1.3	Oral Cavity and Pharynx	7.3	-5.2	Corpus and Uterus, NOS <sup>f</sup>	9.4	0.2
Kidney and Renal Pelvis	7.3	3.6*	Leukemia	6.5	-2.3	Thyroid	8.4	5.1*
Leukemia	7.2	-1.5*	Ovary <sup>fh</sup>	5.9	-1.3	Cervix Uteri <sup>f</sup>	6.1	-5.1*
Ovary <sup>fh</sup>	5.3	-0.2	Thyroid	5.8	-1.6	Ovary <sup>fh</sup>	5.9	-1.8

- <sup>a</sup> Top 15 cancer sites selected based on 2003-2007 age-adjusted rates for the race/ethnic group.
- <sup>b</sup> Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).
- <sup>c</sup> Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).
- <sup>d</sup> Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).  
Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.
- <sup>e</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- <sup>f</sup> Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- <sup>g</sup> The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- <sup>h</sup> IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- <sup>\*</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- <sup>-</sup> The APC is significantly different from zero (p<.05).  
Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.